

MedLexis Portfolio Sample: SEO-Optimized Medical Article

Client Brand: Aorta – Heart Health Platform

Developed By: MedLexis | Premier Medical Writing & Content Agency

Content Focus: Patient Education | Chronic Kidney Disease (CKD) | Dialysis

SEO Metadata Panel:

1. **Title:** *Dialysis Explained: What It Is, How It Works, and When It's Needed*
2. **Primary Keyword –** *Dialysis*
3. **Meta Descriptions:** *Learn what dialysis is, how it works, and when it's needed. Understand types, benefits, and daily life with dialysis.*

About This Sample

This article was **professionally written and medically reviewed by MedLexis** for **Aorta**, a patient-focused health brand dedicated to cardiovascular wellness. It exemplifies our commitment to delivering **accurate, engaging, and SEO-optimized medical content** that informs, empowers, and ranks.

Targeting the high-impact keyword “**dialysis**,” this piece was carefully structured to balance **clinical clarity** with **search engine visibility**. It educates readers on key topics, including **when dialysis is needed, the differences between hemodialysis and peritoneal dialysis**, and **real-life patient considerations**—all presented with empathy, authority, and precision.

As part of our portfolio, this article demonstrates MedLexis’s ability to:

- Translate complex medical concepts into **reader-friendly content**
- Align medical accuracy with **SEO best practices**
- Support health-focused brands with **strategic, high-performing publications**

Whether you're a healthcare organization, publisher, or medical startup, MedLexis delivers content that builds **credibility**, drives **traffic**, and earns **trust**.



Dialysis Explained: What It Is, How It Works, and When It's Needed

The kidneys play a vital role in maintaining the body's internal balance by filtering waste products, excess fluids, and toxins from the bloodstream. They also regulate blood pressure, electrolyte levels, and the production of red blood cells.

When kidney function declines severely—a condition known as **kidney failure** or **end-stage renal disease (ESRD)**—the body can no longer effectively eliminate waste or maintain fluid and chemical balance. Left untreated, this can lead to life-threatening complications.

Dialysis is a life-sustaining treatment that replaces the kidneys' filtering function when it can no longer work adequately. It removes harmful waste and excess fluids, helping to restore the body's balance. For millions of people worldwide living with ESRD, dialysis is not just a treatment—it is a vital bridge to survival and quality of life.

What Is Dialysis?

Dialysis is a medical therapy used to replicate the essential functions of healthy kidneys in individuals with **severe kidney failure**. It involves removing waste products, excess fluids, and toxins from the bloodstream when the kidneys can no longer do so effectively.

When kidney function falls below **10–15% of normal capacity**, dialysis becomes medically necessary to prevent life-threatening complications such as fluid overload, high potassium levels, and uremia. This typically occurs in the advanced stages of **chronic kidney disease (CKD)**, also known as **end-stage renal disease (ESRD)**.



Quick Facts:

- More than **3.5 million people** worldwide rely on dialysis.
 - In the **United States**, over **800,000 people** live with ESRD.
 - Approximately **70%** of U.S. patients with kidney failure are treated with dialysis; others may receive kidney transplants or opt for conservative management.
-

When Is Dialysis Needed?

Dialysis becomes necessary when kidney function declines to **less than 10–15%** of its normal capacity—this is referred to as **end-stage kidney disease (ESKD)**, the most advanced stage of **chronic kidney disease (CKD)**. At this point, the kidneys can no longer effectively remove sufficient waste or excess fluid to maintain the body's balance.

Dialysis may also be needed in **acute kidney injury (AKI)**—a sudden, often reversible loss of kidney function due to illness, injury, or medication effects. In such cases, dialysis is temporary until kidney function improves.

Common Signs You May Need Dialysis:

- Persistent fatigue and weakness
- Swelling in the legs, ankles, or feet (peripheral edema)
- Shortness of breath due to fluid overload
- Nausea, vomiting, or loss of appetite
- Mental confusion or trouble concentrating (uremic encephalopathy)
- Decreased or no urine output

Physicians base the decision to start dialysis not only on **lab values** (e.g., creatinine levels, **estimated glomerular filtration rate [eGFR]**, serum potassium) but also on a person's **clinical symptoms**, nutritional status, and overall well-being. The goal is to initiate treatment **before life-threatening complications** arise.

Types of Dialysis

Dialysis can be performed in several ways, depending on the patient's health, lifestyle, and preferences. The **two primary types** are:

1. Hemodialysis (HD)

Hemodialysis uses a machine and a special filter called a **dialyzer** to clean the blood. Blood is withdrawn from the body, filtered through the machine, and returned via vascular access—often through a fistula or catheter.

- Typically performed in a **dialysis center**
- Sessions occur **3 times per week**, lasting **3–5 hours** each
- Also available as **Home Hemodialysis (HHD)** with appropriate training

Hemodialysis is the most widely used dialysis method worldwide.

2. Peritoneal Dialysis (PD)

In peritoneal dialysis, the **peritoneum**—the natural lining of the abdominal cavity—acts as a filter. A **sterile dialysate** solution is infused through a catheter into the abdomen. After waste and fluids are absorbed, the solution is drained and replaced.

Types include:

- **Continuous Ambulatory Peritoneal Dialysis (CAPD)** – done manually 3–5 times daily
- **Automated Peritoneal Dialysis (APD)** – performed overnight using a machine

Peritoneal dialysis is usually done **at home**, offering more independence and flexibility.

Choosing the Right Type

Each method has unique advantages and considerations. Factors include:

- Medical condition and comorbidities
- Lifestyle and home environment
- Personal preferences
- Support system availability

A **nephrologist** will help guide patients in selecting the most appropriate option based on their needs and goals.

HEMODIALYSIS (HD) VS. PERITONEAL DIALYSIS (PD)

Criteria	Hemodialysis (HD)	Peritoneal Dialysis (PD)
Procedure	Blood is filtered externally through a machine (dialyzer). Requires vascular access (fistula, graft, or catheter).	Uses the body's peritoneal membrane as a natural filter. Requires a permanent abdominal catheter.
Schedule	<ul style="list-style-type: none">• 3-5 hours/session• 3 times/week• In-center or home (with training)	<ul style="list-style-type: none">• CAPD: 3-5 manual exchanges/day• APD: Automated overnight cycles (8-10 hours)
Lifestyle	A fixed schedule; may cause post-treatment fatigue. Travel requires coordination with dialysis centers.	Flexible; can be done at home/work. Easier for travel (portable equipment).
Risks	<ul style="list-style-type: none">• Infection at access site• Low blood pressure during treatment• Blood clotting	<ul style="list-style-type: none">• Peritonitis (abdominal infection)• Hernias (due to fluid pressure)• Catheter-related infections
Best For	Patients needing rapid toxin removal or with: <ul style="list-style-type: none">• Cardiovascular instability• Limited home support	Patients prioritizing: <ul style="list-style-type: none">• Independence• Gentler, continuous clearance• Fewer dietary restrictions

How Dialysis Works: The Science Behind the Process

Dialysis is a life-sustaining therapy that **replicates key functions of the kidneys**, specifically the removal of metabolic waste products, regulation of fluid and electrolyte balance, and maintenance of acid-base homeostasis.

1. Hemodialysis (HD): Artificial Filtration via a Dialyzer

In hemodialysis, the patient's blood is circulated through a machine known as a **dialyzer**—often referred to as an "artificial kidney."

- The dialyzer contains **semi-permeable membranes** that separate the blood from a sterile fluid called **dialysate**.

- Through the processes of **diffusion**, **osmosis**, and **ultrafiltration**, waste products (e.g., urea, creatinine), excess electrolytes (e.g., potassium, phosphate), and water are removed.
- Clean blood is returned to the patient's body via a vascular access point.

2. Peritoneal Dialysis (PD): Natural Filtration via the Peritoneum

In peritoneal dialysis, the body's **peritoneal membrane** acts as the filtering surface.

- A catheter is surgically placed into the abdominal cavity.
- A **dialysate solution** is instilled into the cavity, where it dwells for several hours.
- During this time, waste products and fluid pass across the peritoneal membrane into the dialysate.
- The used fluid is then drained and replaced with a fresh solution.

Efficiency Factors in Both Modalities:

- **Membrane surface area** (e.g., peritoneum or dialyzer membrane)
- **Concentration gradients** of waste and electrolytes
- **Blood flow rate** and **dialysate flow rate**
- **Dwell time** in PD or dialysis duration in HD

End Goal: Restoring Balance

Both dialysis modalities aim to:

- Prevent **uremia** (buildup of toxic waste in the blood)
- Correct **electrolyte disturbances** (e.g., hyperkalemia)
- Manage **fluid overload**
- Stabilize **acid-base balance** (prevent acidosis)

While dialysis cannot fully replace all kidney functions, such as erythropoietin production or vitamin D activation, it effectively performs **critical filtration and fluid regulation**, supporting survival and quality of life in patients with kidney failure.

What to Expect During Dialysis: Process and Experience

Understanding what happens during dialysis can ease anxiety and help patients feel more prepared. While each person's experience may vary slightly depending on the type of dialysis used, the overall process is structured, supervised, and focused on patient safety and comfort.

1. Hemodialysis (HD): Step-by-Step

Hemodialysis is typically performed in a dialysis center three times a week, with each session lasting about 3–5 hours.

- 1. Arrival and Preparation:** Patients are weighed and assessed for vital signs. Access sites (fistula, graft, or catheter) are cleaned and prepared.
- 2. Connection to the Dialysis Machine:** Needles are inserted into the vascular access point, and blood is circulated through the dialyzer.
- 3. The Dialysis Process:** Waste products, extra electrolytes, and fluids are removed from the blood while patients rest, read, or use mobile devices.
- 4. Monitoring:** Nurses continuously monitor blood pressure, heart rate, and machine performance to ensure safety.
- 5. Completion and Recovery:** After treatment, needles are removed, and pressure is applied to stop bleeding. Patients are reweighed and observed briefly before leaving.

2. Peritoneal Dialysis (PD): At-Home Routine

Peritoneal dialysis is typically performed at home. There are two main types:

- 1. Continuous Ambulatory Peritoneal Dialysis (CAPD):** Performed manually 3–5 times daily, each exchange takes about 30–40 minutes.
- 2. Automated Peritoneal Dialysis (APD):** Done overnight using a machine (cycler) while the patient sleeps.

Patient Experience: Physical and Emotional Considerations

- **Fatigue** is common, especially after hemodialysis sessions.
- **Cramps or low blood pressure** may occur during or after dialysis.
- Emotional challenges such as stress, anxiety, or frustration are not uncommon.

Risks, Complications, and Common Myths

Dialysis is a life-saving treatment, but like any medical intervention, it carries potential risks. Understanding these helps patients make informed decisions and manage their health effectively.

Common Risks and Complications

1. Hemodialysis Risks:

- **Infection** at the vascular access site (especially with catheters)
- **Low blood pressure (hypotension)** during or after treatment
- **Muscle cramps** or nausea
- **Access failure** (e.g., clotting or narrowing of the fistula/graft)
- **Fatigue or dizziness** following sessions

2. Peritoneal Dialysis Risks:

- **Peritonitis:** A serious infection of the abdominal lining
- **Catheter site infections**
- **Hernias** from increased abdominal pressure
- **Dialysate leakage** around the catheter

Most complications can be minimized with **proper training, hygiene, and close monitoring** by healthcare professionals.

Life on Dialysis — Lifestyle Changes, Diet, and Support Systems

Dialysis does more than replace kidney function — it reshapes daily life. Adapting to this change is essential for optimizing health, preserving independence, and maintaining emotional well-being.

1. Lifestyle Adjustments

Living with dialysis requires thoughtful planning:

- **Time Commitment:** In-center hemodialysis typically takes 9–15 hours per week; peritoneal dialysis must be integrated into daily or nightly routines.
- **Fatigue Management:** Patients often need rest after treatments. Prioritizing sleep, stress reduction, and moderate physical activity can help improve energy.
- **Work and Travel:** Many patients continue working, especially with home-based dialysis. Portable equipment and dialysis networks support travel for both HD and PD users.

2. Renal-Friendly Diet

Kidney disease alters how the body handles fluids, minerals, and waste. A dialysis-appropriate diet is crucial to avoid complications like fluid overload or electrolyte imbalances.

- **Limit Sodium and Potassium:** Excess levels can lead to hypertension or heart arrhythmias.
- **Control Phosphorus Intake:** High phosphorus can weaken bones and damage blood vessels.
- **Protein Needs Increase:** Unlike pre-dialysis, patients on dialysis may need more high-quality protein.
- **Fluid Restrictions:** Tailored to each patient, based on urine output and dialysis type.

A registered dietitian specializing in renal care can personalize meal plans to balance nutrition with medical needs.

3. Emotional and Psychological Support

Chronic illness impacts mental health. Common experiences include:

- Depression or anxiety
- Body image concerns (e.g., due to access sites or fatigue)
- Social withdrawal

Support resources include:

- Counseling and psychotherapy
- Peer support groups (online or in-person)
- Education programs to build confidence in managing dialysis

The Importance of a Strong Support System

Family, caregivers, and healthcare professionals form a vital support network. Successful dialysis often depends on:

- Reliable transportation to appointments
- Adherence to medication and fluid restrictions
- Regular communication with the care team

Empowerment through education, routine, and connection helps patients not just survive, but thrive on dialysis.

Alternatives to Long-Term Dialysis — Kidney Transplant and Conservative Care

Dialysis is not the only path forward for patients with kidney failure. Two primary alternatives — kidney transplantation and conservative (non-dialysis) care — may be more appropriate depending on individual health, goals, and preferences.

Kidney Transplantation: The Gold Standard

A kidney transplant offers the greatest potential for freedom from dialysis and improved quality of life.

Benefits include:

- More natural kidney function
- Fewer dietary restrictions
- Greater energy and life expectancy
- No need for ongoing dialysis treatments

Eligibility Considerations:

- Good overall health
- No active infections or untreated cancers
- Commitment to lifelong immunosuppressive therapy

Types of Transplants:

- **Living donor:** Often preferred due to better outcomes and shorter wait times.
- **Deceased donor:** Based on organ availability and match.

Transplant recipients require **lifelong follow-up** and **strict adherence to medications** to prevent organ rejection.

Conservative Management (Supportive Care Without Dialysis)

For some individuals, especially older adults or those with significant comorbidities, dialysis may not improve their quality of life or align with personal values.

Conservative kidney management focuses on:

- **Symptom relief** (fatigue, nausea, shortness of breath)
Blood pressure and anemia control
- **Nutritional support**
- **Palliative and hospice care as needed**

This approach emphasizes **comfort, dignity, and quality of life** over life extension. It is a valid, compassionate option when dialysis is not in the patient's best interest.

Informed Choice Matters

Choosing between dialysis, transplant, or conservative care is deeply personal. Shared decision-making with nephrologists, families, and palliative care teams ensures care aligns with the patient's goals and values.

Understanding Dialysis and Taking Charge of Your Kidney Health

Dialysis is a life-saving therapy that supports individuals with kidney failure when the kidneys can no longer perform their essential functions. Whether through hemodialysis or peritoneal dialysis, this treatment helps manage waste, fluids, and toxins, making it possible for patients to live longer and maintain quality of life.

While the journey may come with challenges, understanding your options, such as transplantation or conservative care, empowers you to make informed, meaningful decisions in partnership with your healthcare team. With the right medical support, lifestyle adjustments, and emotional resources, it is possible to live fully and purposefully on dialysis or beyond.

From all of us at **AORTA**, we extend our support and best wishes to you and your loved ones navigating kidney health. We believe that knowledge is power, and by staying informed, you are taking a crucial step toward better well-being.

Explore more on AORTA:

Looking for more trustworthy insights on kidney and heart health? Visit our blog for related articles on chronic kidney disease, the heart-kidney connection, and nutrition tips for dialysis patients.

Share This Article:

If you found this guide helpful, please share it with friends, family, or on social media. Help us spread awareness and empower others to take control of their health.

References

1. National Institute of Diabetes and Digestive and Kidney Diseases. Dialysis. US Department of Health and Human Services; 2023. Accessed May 28, 2025. <https://www.niddk.nih.gov/health-information/kidney-disease/kidney-failure/dialysis>
2. Centers for Disease Control and Prevention. Chronic kidney disease in the United States, 2023. US Department of Health and Human Services; 2023. Accessed May 28, 2025. <https://www.cdc.gov/kidneydisease/publications-resources/ckd-national-facts.html>
3. About dialysis. National Kidney Foundation. Accessed May 28, 2025. <https://www.kidney.org/atoz/content/dialysisinfo>
4. Dialysis: what you can expect. Mayo Clinic. Published October 12, 2023. Accessed May 28, 2025. <https://www.mayoclinic.org/tests-procedures/dialysis/about/pac-20384827>
5. Kidney Disease: Improving Global Outcomes (KDIGO). KDIGO 2020 clinical practice guidelines for chronic kidney disease. *Kidney Int.* 2020;98(4)(suppl):S1-S115. doi:10.1016/j.kint.2020.06.005
6. United States Renal Data System. 2023 USRDS annual data report: epidemiology of kidney disease in the United States. National Institutes of Health; 2023. Accessed May 28, 2025. <https://adr.usrds.org/2023>
7. Dialysis overview. National Health Service (NHS). Updated February 10, 2023. Accessed May 28, 2025. <https://www.nhs.uk/conditions/dialysis/>
8. Levin A, Tonelli M, Bonventre JV, et al. Global kidney health 2017 and beyond: a roadmap for closing gaps in care, research, and policy. *Lancet.* 2017;390(10105):1888-1917. doi:10.1016/S0140-6736(17)30788-2
9. Jha V, Garcia-Garcia G, Iseki K, et al. Chronic kidney disease: global dimension and perspectives. *Lancet.* 2013;382(9888):260-272. doi:10.1016/S0140-6736(13)60687-X
10. Conservative care guidelines and recommendations. Canadian Society of Nephrology. Accessed May 28, 2025. <https://www.csnsn.ca/>

MedLexis Portfolio Disclaimer

This portfolio sample is intended solely for educational and promotional purposes. It does not constitute medical advice, diagnosis, or treatment, nor does it represent the views or content of any actual client unless explicitly stated. All medical content is based on publicly available, evidence-based information and created to demonstrate the writing expertise of MedLexis. For real-world applications, always consult a qualified medical professional. MedLexis upholds the highest standards of client confidentiality, content ethics, and scientific accuracy.

Let's Elevate Your Medical Communication.

 Contact us at Contact@Medicallexis.com |  Visit: www.Medicallexis.com

Contact Details

Writer Name	Writer Email	MedLexis Contact Email
Dr. Hayder Mazin	Hayder@Medicallexis.com	Contact@Medicallexis.com



[© 2025 MedLexis. All rights reserved.](#)

[Precision in Writing, Trust in Health.](#)